

Due to regulatory changes, the content of the following Patient Information Leaflet may vary from the one found in your medicine pack. Please compare the 'Leaflet prepared/revised date' towards the end of the leaflet to establish if there have been any changes.

If you have any doubts or queries about your medication, please contact your doctor or pharmacist.



Package Leaflet: Information for the user

Levitra 10 mg orodispersible tablets Vardenafil

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Levitra is and what it is used for
2. What you need to know before you take Levitra
3. How to take Levitra
4. Possible side effects
5. How to store Levitra
6. Contents of the pack and other information

1. What Levitra is and what it is used for

Levitra contains vardenafil, a member of a class of medicines called phosphodiesterase type 5 inhibitors. They are used for the treatment of erectile dysfunction in adult men, a condition which implies difficulties in getting or keeping an erection.

At least one in ten men has trouble getting or keeping an erection at some time. There may be physical or psychological causes, or a mixture of both. Whatever the cause is, due to muscle and blood vessel changes not enough blood stays in the penis to make it hard and keep it hard.

Levitra will only work when you are sexually stimulated. It reduces the action of the natural chemical in your body which makes erections go away. Levitra allows an erection to last long enough for you to satisfactorily complete sexual activity.

2. What you need to know before you take Levitra

Do not take Levitra

- If you are allergic to vardenafil or any of the other ingredients of this medicine (listed in section 6). Signs of an allergic reaction include a rash, itching, swollen face or lips and shortness of breath.
- If you are taking medicines containing nitrates, such as glycerol trinitrate for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure.
- If you are taking ritonavir or indinavir, medicines used to treat human immunodeficiency virus (HIV) infections.
- If you are over 75 years of age and are taking ketoconazole or itraconazole, anti-fungal medicines.
- If you have a severe heart or liver problem.
- If you are having kidney dialysis.
- If you have recently had a stroke or heart attack.
- If you have or have had low blood pressure.
- If your family has a history of degenerative eye diseases (such as retinitis pigmentosa).
- If you have ever had a condition involving loss of vision due to damage to the optic nerve from insufficient blood supply known as non-arteritic ischemic optic neuropathy (NAION).

Warnings and precautions

Talk to your doctor or pharmacist before taking Levitra.

Take special care with Levitra

- If you have heart trouble. It may be risky for you to have sex.
- If you suffer from irregular heart beat (cardiac arrhythmia) or inherited heart diseases affecting your electrocardiogram.
- If you have a physical condition affecting the shape of the penis. This includes conditions called angulation, Peyronie's disease and cavernosal fibrosis.
- If you have an illness that can cause erections which won't go away (priapism). These include sickle cell disease, multiple myeloma and leukaemia.
- If you have stomach ulcers (also called gastric or peptic ulcers).
- If you have a bleeding disorder (such as haemophilia).
- If you are using any other treatments for erection difficulties, including Levitra film-coated tablets (see section: Other medicines and Levitra).
- If you experience sudden decrease or loss of vision, stop taking Levitra and contact your doctor immediately.

Children and adolescents

Levitra is not intended for use by children or adolescents under 18.

Other medicines and Levitra

Please tell your doctor or pharmacist if you are using, have recently used or might use any other medicines, including medicines obtained without a prescription.

Some medicines may cause problems, especially these:

- Nitrates, medicines for angina, or nitric oxide donors, such as amyl nitrite. Taking these medicines with Levitra could seriously affect your blood pressure. Talk to a doctor without taking Levitra.
- Medicine for the treatment of arrhythmias, such as quinidine, procainamide, amiodarone or sotalol.
- Ritonavir or indinavir, medicines for HIV. Talk to a doctor without taking Levitra.
- Ketoconazole or itraconazole, anti-fungal medicines.
- Erythromycin, or clarithromycin, macrolide antibiotics.
- Alpha-blockers, a type of medicine used to treat high blood pressure and enlargement of the prostate (as benign prostatic hyperplasia).

Do not use Levitra orodispersible tablets combined with any other treatment for erectile dysfunction, including Levitra film-coated tablets.

Levitra with food and drink

- You can take Levitra orodispersible tablets with or without food, but do not take this medicine with any liquid.
- Don't drink grapefruit juice when you use Levitra. It can interfere with the usual effect of the medicine.
- Alcoholic drink can make erection difficulties worse.

Pregnancy and breast-feeding

Levitra is not for use by women.

Driving and using machines

Levitra might make some people feel dizzy or affect their vision. If you feel dizzy, or if your vision is affected after taking Levitra don't drive or operate any tools or machines.

Levitra 10 mg orodispersible tablets contains Aspartame and Sorbitol.


- Aspartame is a source of phenylalanine, and may be harmful for people with a disorder called phenylketonuria.
- Sorbitol: If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Levitra orodispersible tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. The recommended dose is 10 mg.

Take Levitra tablet about 25 to 60 minutes before sexual activity. With sexual stimulation you may achieve an erection anywhere from 25 minutes up to four to five hours after taking Levitra.

- Do not remove the orodispersible tablet from the blister until you are going to take it. With dry hands, pull the blister edge from the blister holder and press gently on top to release the tablet on your hand. Do not crush the tablet.
- Place the entire orodispersible tablet in the mouth, on the tongue, where it will dissolve in seconds, then swallow with saliva. The orodispersible tablet must be taken without any liquid.



Do not take Levitra orodispersible tablets with any other forms of Levitra.

Do not take Levitra more than once a day.

Tell your doctor if you think Levitra is too strong or too weak. He or she may suggest a switch to an alternative Levitra formulation with a different dose, depending on how well it works for you.

If you take more Levitra than you should

Men who take too much Levitra may experience more side effects or may get severe back pain. If you take more Levitra than you should, tell your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the effects are mild or moderate.

Partial, sudden, temporary or permanent decrease or loss of vision in one or both eyes has been experienced by patients. Stop taking Levitra and contact your doctor immediately.
Sudden decrease or loss of hearing has been reported.

The chance of having a side effect is described by the following categories:

Very common:

may affect more than 1 in 10 user

- Headache

Common:

may affect up to 1 in 10 users

- Dizziness
- Flushing
- Blocked or runny nose
- Indigestion

Uncommon:

may affect up to 1 in 100 users

- Swelling of skin and mucous tissue including swollen face, lips or throat.
- Sleep disorder
- Numbness and impaired perception of touch
- Sleepiness
- Effects on vision; redness of the eye, effects on colour vision, eye pain and discomfort, light sensitivity
- Ringing in the ears; vertigo
- Fast heart beat or pounding heart
- Breathlessness
- Stuffy nose
- Acid reflux, gastritis, abdominal pain, diarrhoea, vomiting; feeling sick (*nausea*), dry mouth
- Raised levels of liver enzymes in your blood
- Rash, reddened skin
- Back or muscle pain; increase in blood of a muscle enzyme (*creatine phosphokinase*), muscle stiffness
- Prolonged erections
- Malaise

Rare:

may affect up to 1 in 1,000 users

- Inflammation of the eyes (*conjunctivitis*)
- Allergic reaction
- Anxiety
- Fainting
- Amnesia
- Seizure
- Increased pressure in the eye (*glaucoma*), lacrimation increased
- Effects on the heart (such as heart attack, altered heart beat or angina)
- High or low blood pressure
- Nose bleed
- Effect on results of blood tests to check liver function.

- Sensitivity of the skin to sun light
- Painful erections
- Chest pain

Very rare or not known:

may affect less than 1 in 10,000 users or frequency cannot be estimated from the available data

- Blood in the urine (*Haematuria*)
- Penile bleeding (*Penile Haemorrhage*)
- Presence of blood in the semen (*Haematospermia*)

Reporting of side effects

If you get any side effects, talk to your doctor. You can also report side effects directly (see details below). By reporting side effects, you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme

Website: www.mhra.gov.uk/yellowcard

Ireland

IMB Pharmacovigilance

Earlsfort Terrace

IRL - Dublin 2

Tel: +353 1 6764971

Fax: +353 1 6762517

Website: www.imb.ie

e-mail: imbpharmacovigilance@imb.ie

Malta

ADR Reporting

The Medicines Authority

Post-Licensing Directorate

203 Level 3, Rue D'Argens

GŻR-1368 Gżira

Website: www.medicinesauthority.gov.mt

e-mail: postlicensing.medicinesauthority@gov.mt

5. How to store Levitra

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after “EXP”. The expiry date refers to the last day of that month.

Store in the original package in order to protect from humidity and light.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Levitra orodispersible tablets contains

- The active substance is vardenafil. Each orodispersible tablet contains 10 mg of vardenafil (as hydrochloride).

- The other ingredients of the tablets are:
Magnesium stearate, aspartame (E951), peppermint flavour, mannitol (E421) sorbitol (E420), crospovidone and silica colloidal hydrated.

What Levitra 10 mg orodispersible tablets looks like and contents of the pack

Levitra 10 mg orodispersible tablets are round and white. They are provided in packs of 1, 2, 4 or 8 tablets (containing Alu/Alu blisters or Alu/Alu blisters in blister holders).

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Bayer Pharma AG

13342 Berlin

Germany

Manufacturer

Bayer Pharma AG

51368 Leverkusen

Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Bayer SA-NV
Tél/Tel: +32-(0)2-535 63 11

България

Байер България ЕООД
Тел.: +359-(0)2 81 401 01

Česká republika

Bayer s.r.o.
Tel: +420 266 101 111

Danmark

Bayer A/S
Tlf: +45 45 23 50 00

Deutschland

Jenapharm GmbH & Co. KG
Tel: +49-(0)3641-64 8888

Eesti

Bayer OÜ
Tel: +372 655 85 65

Ελλάδα

Bayer Ελλάς ABEE
Τηλ: +30 210 61 87 500

España

Bayer Hispania S.L.
Tel: +34-93-495 65 00

France

Bayer HealthCare
Tél (N° vert): +33-(0)800 87 54 54

Hrvatska

Bayer d.o.o.
Tel: + 385-(0)1-6599 900

Ireland

Bayer Limited
Tel: +353-1-2999313

Ísland

Icepharma hf.
Sími: +354-540 8000

Italia

Bayer S.p.A.
Tel: +39-02-397 81

Κύπρος

NOVAGEM Limited
Τηλ: +357 22 48 38 58

Latvija

SIA Bayer
Tel: +371 67 845 563

Lietuva

UAB Bayer
Tel: +37 05 23 36 868

Luxembourg/Luxemburg

Bayer SA-NV
Tél/Tel: +32-(0)2-535 63 11

Magyarország

Bayer Hungária KFT
Tel.: +36-1-487-4100

Malta

Alfred Gera and Sons Ltd.
Tel: +35 621 44 62 05

Nederland

Bayer B.V.
Tel: +31-(0)297-28 06 66

Norge

Bayer AS
Tlf: +47-24 11 18 00

Österreich

Bayer Austria Ges. m. b. H.
Tel: +43-(0)1-711 460

Polska

Bayer Sp. z o.o.
Tel.: +48-22-572 35 00

Portugal

Bayer Portugal S.A.
Tel: +351-21-416 42 00

România

SC Bayer SRL.
Tel: +40-21-529 59 00

Slovenija

Bayer d. o. o.
Tel: +386-1-58 14 400

Slovenská republika

Bayer, spol. s r.o.
Tel: +421 2-59 21 31 11

Suomi/Finland

Bayer Oy
Puh/Tel: +358-20-785 21

Sverige

Bayer AB
Tel: +46-(0)8-580 223 00

United Kingdom

Bayer plc
Tel: +44-(0)1635-56 3000

This leaflet was last revised in 04/2014.

Detailed information on this medicine is available on the European Medicines Agency web site:

<http://www.ema.europa.eu>

